

## 4.1 Graphical resolution and standard form

Given the problem

$$\begin{aligned} \min \quad & \underline{c}^T \underline{x} \\ & A\underline{x} \geq \underline{b} \\ & \underline{x} \geq \underline{0} \end{aligned}$$

where

$$\underline{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \quad \underline{c} = \begin{pmatrix} 16 \\ 25 \end{pmatrix}, \quad \underline{b} = \begin{pmatrix} 4 \\ 5 \\ 9 \end{pmatrix}, \quad A = \begin{pmatrix} 1 & 7 \\ 1 & 5 \\ 2 & 3 \end{pmatrix}$$

1. Solve it graphically.
2. Restate it in standard form, identify  $B$ ,  $N$ , and the corresponding partition of the cost vector w.r.t. the vertex which is optimal for the related polyhedron.

## 4.2 Geometry of linear programming

Given the linear program

$$\begin{aligned} \max \quad z = \quad & 3x_1 + 2x_2 \\ & 2x_1 + x_2 \leq 4 & (1) \\ & -2x_1 + x_2 \leq 2 & (2) \\ & x_1 - x_2 \leq 1 & (3) \\ & x_1, x_2 \geq 0 \end{aligned}$$

1. Solve it graphically, indicating, for each variable and for the objective function, the value that it takes in an optimal solution.
2. Determine the basic solutions related to all the vertices of the polyhedron of the feasible region.
3. Indicate the sequence of basic solutions that are visited by the simplex algorithm (let  $x_1$  be the first variable to enter the basis).
4. Determine the reduced costs for the basic solution associated to the vertices  $((\text{eq. 1}) \cap (\text{eq. 2}))$  and  $((\text{eq. 1}) \cap (\text{eq. 3}))$ , where (eq.  $i$ ) is obtained from (i), by substituting  $=$  for  $\leq$ .
5. Show, geometrically, that the gradient of the objective function is a conic combination (i.e. a nonnegative linear combination) of the gradients of the constraints which are active at an optimal vertex. Indicate the value taken by the objective function in that vertex. *Note:* All the constraints must be in  $\leq$  form, since the problem is a maximization one (e.g.,  $x_1 \geq 0$  must be rewritten as  $-x_1 \leq 0$ ).

6. Determine the range of values for the right hand side  $b_1$  of constraint (1) for which the optimality of the basis solution is preserved.
7. Indicate for which values of the objective function coefficients  $((x_1 = 0) \cap (\text{eq. 2}))$  is an optimal vertex.
8. Determine the range of values for the right hand side  $b_2$  of constraint (2) for which the feasible region is (a) empty (b) contains a single point.
9. Indicate the values for  $c_1$  for which there are multiple optimal solutions.

### 4.3 Simplex algorithm with Bland's rule

Given the linear program

$$\begin{aligned} \min \quad z &= x_1 - 2x_2 \\ 2x_1 &\quad + 3x_3 = 1 \\ 3x_1 + 2x_2 &- x_3 = 5 \\ x_1, x_2, x_3 &\geq 0 \end{aligned}$$

solve it via the two phases simplex algorithm applying Bland's rule.

## SOLUTION

## 4.1 Graphical resolution and standard form

1. The equations associated to the system are  $A\underline{x} = \underline{b}$

$$x_1 + 7x_2 = 4 \quad (4)$$

$$x_1 + 5x_2 = 5 \quad (5)$$

$$2x_1 + 3x_2 = 9 \quad (6)$$

The corresponding lines are shown in Figure 1. The level curves of the objective function




Figure 1: The polyhedron is unbounded

are  $z = 16x_1 + 25x_2$ , or, equivalently,  $x_2 = -\frac{16}{25}x_1 + \frac{z}{25}$ . Inspecting Figure 2, we observe that vertex  $R$  is the unique optimal solution. Since vertex  $R$  is the intersection of equations (5) and (6), we obtain it by solving

$$\begin{aligned} x_1 + 5x_2 &= 5 \\ 2x_1 + 3x_2 &= 9 \end{aligned}$$

obtaining  $x_1 = \frac{30}{7}, x_2 = \frac{1}{7}$ .

2. To express the problem in standard form, we introduce 3 slack variables,  $s_1, s_2, s_3$ , one per




Figure 2: Graphical resolution:  $R$  is an optimal solution

constraint. The new problem is

$$\begin{aligned} \min \quad & \underline{c}'^T \underline{x}' \\ & A' \underline{x}' = \underline{b} \\ & \underline{x}' \geq \underline{0} \end{aligned}$$

where  $\underline{x}' = (x_1, x_2, s_1, s_2, s_3)^T$ ,  $\underline{c}' = (16, 25, 0, 0, 0)$ , and

$$A' = \left( \begin{array}{cc|ccc} 1 & 7 & -1 & 0 & 0 \\ 1 & 5 & 0 & -1 & 0 \\ 2 & 3 & 0 & 0 & -1 \end{array} \right) = (A| -I).$$

Since  $R$  is obtained by intersecting (4) and (5), the slack variables of the corresponding constraints are null in  $R$ . By letting  $\underline{x}_B = (x_1, x_2, s_1)$  and  $\underline{x}_N = (s_2, s_3)$ , we partition the variables into  $\underline{x}' = (\underline{x}_B | \underline{x}_N)$ . The corresponding partition of matrix  $A'$  is

$$A' = \left( \begin{array}{ccc|cc} 1 & 7 & -1 & 0 & 0 \\ 1 & 5 & 0 & -1 & 0 \\ 2 & 3 & 0 & 0 & -1 \end{array} \right) = (B|N).$$

The basic variables in  $R$  have values  $\underline{x}_B = B^{-1}\underline{b}$ . Therefore,

$$B^{-1} = \frac{1}{7} \begin{pmatrix} 0 & -3 & 5 \\ 0 & 2 & -1 \\ -7 & 11 & -2 \end{pmatrix}$$

and  $\underline{x}_B = (\frac{30}{7}, \frac{1}{7}, \frac{9}{7})$ , where  $\underline{x}_B = (x_1, x_2)'$ .

## 4.2 Geometry of linear programming

1. The equations associated to constraints (1), (2), (3) are

$$2x_1 + x_2 = 4 \quad (\text{eq. 1})$$

$$-2x_1 + x_2 = 2 \quad (\text{eq. 2})$$

$$x_1 - x_2 = 1 \quad (\text{eq. 3})$$

or, equivalently,

$$x_2 = -2x_1 + 4$$

$$x_2 = 2x_1 + 2$$

$$x_2 = x_1 - 1.$$

The level curves of the objective function are  $z = 3x_1 + 2x_2$ , i.e.,  $x_2 = -\frac{3x_1}{2} + \frac{z}{2}$ . The polyhedron  $PQRS$  of the feasible solutions is shown in Figure 3. The unique optimal solution is achieved at vertex  $P = (\frac{1}{2}, 3)$ , where  $z^* = \frac{15}{2}$ .




Figure 3: Polyhedron of the feasible solutions

2. To express the problem in standard form, we introduce 3 slack variables,  $s_1, s_2, s_3$ , one per constraint. The problem is

$$\begin{aligned} \max \quad & z = \underline{c}'^T \underline{x}' \\ & A' \underline{x}' = \underline{b} \\ & \underline{x}' \geq \underline{0}, \end{aligned}$$

where

$$\underline{x}' = \begin{pmatrix} x_1 \\ x_2 \\ s_1 \\ s_2 \\ s_3 \end{pmatrix}, \quad \underline{c}' = \begin{pmatrix} 3 \\ 2 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad \underline{b}' = \underline{b}, \quad A' = (A|I) = \left( \begin{array}{cc|ccc} 2 & 1 & 1 & 0 & 0 \\ -2 & 1 & 0 & 1 & 0 \\ 1 & -1 & 0 & 0 & 1 \end{array} \right).$$

- (a) Vertex  $P$ :  $s_1 = 0, s_2 = 0$ , where  $\underline{x}_B = (x_1, x_2, s_3)$ ,  $\underline{x}_N = (s_1, s_2)$ ,

$$B = \begin{pmatrix} 2 & 1 & 0 \\ -2 & 1 & 0 \\ 1 & -1 & 1 \end{pmatrix}, \quad N = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

- (b) Vertex  $Q$ :  $s_1 = 0, s_3 = 0$ , where  $\underline{x}_B = (x_1, x_2, s_2)$ ,  $\underline{x}_N = (s_1, s_3)$ ,

$$B = \begin{pmatrix} 2 & 1 & 0 \\ -2 & 1 & 1 \\ 1 & -1 & 0 \end{pmatrix}, \quad N = \begin{pmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

- (c) Vertex  $R$ :  $x_2 = 0, s_3 = 0$ , where  $\underline{x}_B = (x_1, s_1, s_2)$ ,  $\underline{x}_N = (x_2, s_3)$ ,

$$B = \begin{pmatrix} 2 & 1 & 0 \\ -2 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad N = \begin{pmatrix} 1 & 0 \\ 1 & 0 \\ -1 & 1 \end{pmatrix}.$$

- (d) Vertex  $O$ :  $x_1 = 0, x_2 = 0$ , where  $\underline{x}_B = (s_1, s_2, s_3)$ ,  $\underline{x}_N = (x_1, x_2)$ ,

$$B = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad N = \begin{pmatrix} 2 & 1 \\ -2 & 1 \\ 1 & -1 \end{pmatrix}.$$

- (e) Vertex  $S$ :  $x_1 = 0, s_2 = 0$ , where  $\underline{x}_B = (x_2, s_1, s_3)$ ,  $\underline{x}_N = (x_1, s_2)$ ,

$$B = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 0 \\ -1 & 0 & 1 \end{pmatrix}, \quad N = \begin{pmatrix} 2 & 0 \\ -2 & 1 \\ 1 & 0 \end{pmatrix}.$$

3. Assume that, initially, the basic solution is given by the slack variables, i.e.,  $\underline{x}_B = (s_1, s_2, s_3)$ . Variables  $x_1, x_2$  are nonbasic. The solution corresponds to vertex  $O$  (the origin). Assuming that  $x_1$  becomes basic, we are increasing the value of  $x_1$ , i.e., we are

moving on segment  $O - R$ . Variables  $s_1$  and  $s_3$  decrease as  $x_1$  increases. The first one to become 0 is  $s_3$ . We reach vertex  $R$ , where  $\underline{x}_B = (x_1, s_1, s_2)$ . The next variable to become basic is  $x_2$ . We are moving on segment  $R - Q$ , and obtain the next solution in  $Q$ , where  $\underline{x}_B = (x_1, x_2, s_2)$ . Then  $s_3$  becomes basic and  $s_2$  becomes nonbasic, i.e., we move on segment  $Q - P$ , reaching the unique optimal vertex  $P$ , where  $\underline{x}_B = (x_1, x_2, s_3)$ ,

4. The vertex given by (1)  $\cap$  (2) is  $P$ . That given by (1)  $\cap$  (3) is  $Q$ . The reduced costs are  $\bar{c} = \underline{c}^T - \underline{c}_B^T B^{-1}A$ . Observe that  $\bar{c}$  is zero for basic variables, and (possibly!) nonzero for nonbasic ones. Therefore, we only consider  $\bar{c}_N = \underline{c}_N^T - \underline{c}_B^T B^{-1}N$ . In vertex  $P$ , we have  $B$  and  $N$  as in (b)i, so

$$B^{-1}N = \frac{1}{4} \begin{pmatrix} 1 & -1 \\ 2 & 2 \\ 1 & 3 \end{pmatrix}$$

where  $\underline{c}'_B = (3, 2, 0)$  and  $\underline{c}'_N = (0, 0)$ . We obtain  $\bar{c}_N = (-\frac{7}{4}, -\frac{1}{4})$ . Since both values are  $\geq 0$  and the problem is maximization one, the basic solution corresponding to vertex  $P$  is optimal. In vertex  $Q$ , we have  $B, N$  as in 4.2(b)ii, so

$$B^{-1}N = \frac{1}{3} \begin{pmatrix} 1 & 1 \\ 1 & -2 \\ 1 & 4 \end{pmatrix}$$

where  $\underline{c}'_B = (3, 2, 0)$  and  $\bar{c}_N = (-\frac{5}{3}, \frac{1}{3})$ . Therefore,  $Q$  is not an optimal solution.

5. The gradient of the objective function is a conic combination of the gradients of the active constraints only in an optimal vertex. It means that any improving direction is infeasible, which implies that the current vertex is optimal. Consider  $P = (\frac{1}{2}, 3)$ . The gradient of the objective function is  $\nabla f = (3, 2)$ . The active constraints, in  $P$ , are (1) and (2). The gradients are  $(2, 1)$  and  $(-2, 1)$ . We are to check whether the system

$$\lambda_1 \begin{pmatrix} 2 \\ 1 \end{pmatrix} + \lambda_2 \begin{pmatrix} -2 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$$


admits a solution where  $\lambda_1 \geq 0$  and  $\lambda_2 \geq 0$ . Since it is of full rank, the solution is unique. Since it amounts to  $\lambda_1 = \frac{7}{4}, \lambda_2 = \frac{1}{4}$ ,  $P$  satisfies the condition. See Figure 4.

We verify that the condition is not satisfied in the nonoptimal vertices  $Q, R, O, S$ .

- Vertex  $Q$ . Active constraints (1), (3) with gradients  $(2, 1), (1, -1)$ . We obtain  $\lambda_1 = \frac{5}{3}, \lambda_2 = -\frac{1}{3} < 0$ . The condition is not satisfied.
- Vertex  $R$ . Active constraints (3),  $-x_2 \leq 0$  with gradients  $(1, -1), (0, -1)$ . We obtain  $\lambda_1 = 3, \lambda_2 = -5 < 0$ . The condition is not satisfied.
- Vertex  $O$ . Active constraints  $-x_1 \leq 0, -x_2 \leq 0$  with gradients  $(-1, 0), (0, -1)$ . We obtain  $\lambda_1 = -3 < 0, \lambda_2 = -2 < 0$ . The condition is not satisfied.
- Vertex  $S$ . Active constraints  $-x_1 \leq 0, (2)$  with gradients  $(-1, 0), (-2, 1)$ . We obtain  $\lambda_1 = -7 < 0, \lambda_2 = 2$ . The condition is not satisfied.

6. *Geometrical solution, without sensitivity analysis.*

For  $b_1 \rightarrow \infty$ , the optimality of the optimal basis is preserved. Let  $S = (0, s)$ . If  $b_1$  decreases, while  $b_1 > s$ , the optimality of the basis is preserved. If  $b_1 = s$ , we have a

Figure 4: Optimality of vertex  $P$ 

degenerate solution which can be expressed both as  $x_B = (x_1, s_3, x_2)$  or  $x_B = (x_1, s_3, s_2)$ . For  $0 < b_1 < s$ ,  $x_1 = 0$  becomes nonbasic and  $s_2$  becomes basic, since the corresponding constraint (2) is no more active. For  $b_1 = 0$ , the problem has a single feasible point,  $(0, 0)$ , and for  $b_1 < 0$  the feasible region is empty.

7.  $S$  is an optimal solution for any objective function with a gradient which is a conic combination of  $(-1, 0)^T$  and  $(-2, 1)^T$ .
8. The feasible region contains a single point,  $Q$ , for  $b_2$  such that  $Q$  is the intersection of the three lines corresponding to the three constraints, i.e., for  $b_2 = -\frac{8}{3}$ . The feasible region is empty for  $b_2 < -\frac{8}{3}$ .
9. We have multiple optimal solutions if the gradient of the objective function is parallel to that of an inequality defining a facet  $f$  of the polyhedron, i.e., if the two gradients are equivalent up to a positive multiplicative factor.

### 4.3 Simplex method with Bland's rule

The problem is already in standard form. Constraints (1)-(3) equal to the system  $A\underline{x} = \underline{b}$ , where  $\underline{x} = (x_1, x_2, x_3)^T$ ,  $\underline{b} = (1, 5)^T$  and

$$A = \begin{pmatrix} 2 & 0 & 3 \\ 3 & 2 & -1 \end{pmatrix}.$$

Since a feasible basic solution is not evident, we apply the two phases simplex method.

#### PHASE I



This phase finds a feasible basic solution, if any. We solve the auxiliary problem

$$\begin{aligned} \min \quad & v = \sum_i y_i \\ \text{s.t.} \quad & A\underline{x} + I\underline{y} = \underline{b} \\ & \underline{x} \geq \underline{0} \\ & \underline{y} \geq \underline{0} \end{aligned}$$

where an artificial  $y_i$  is introduced for each equality constraint. Observe that vector  $\underline{b}$  must be nonnegative. If it is not, we need to multiply the lines where the entry is negative by  $-1$ .

A feasible basic solution for the auxiliary problem is given by  $\underline{x}_B = (y_1, y_2)$ , where the objective function has a strictly positive value. The problem admits an optimal solution of value 0 if and only if the original problem admits a feasible solution. Indeed, to achieve a value of 0, all variables  $y_i$  must be zero, which implies that  $Ax = b$  can be satisfied.

If no optimal solution of value 0 is found, the original problem is infeasible and the algorithm stops.

For the current problem, the auxiliary problem reads  $\min\{y_1 + y_2 \mid \bar{A}\bar{x} = b, x \geq 0, y \geq 0\}$ . The initial basic solution is  $\bar{x}_B = (y_1, y_2)$ . We express the basic variables w.r.t. the nonbasic ones

$$\begin{aligned} y_1 &= 1 - 2x_1 - 3x_3 \\ y_2 &= 5 - 3x_1 - 2x_2 + x_3 \end{aligned}$$

so that the objective function becomes  $v = y_1 + y_2 = 6 - 5x_1 - 2x_2 - 2x_3$ . The initial tableau is

		$x_1$	$x_2$	$x_3$	$y_1$	$y_2$
	-6	-5	-2	-2	0	0
$y_1$	1	2	0	3	1	0
$y_2$	5	3	2	-1	0	1

The reduced costs of the nonbasic variables  $\bar{x}_N = (x_1, x_2, x_3)$  are all negative. Using Bland's rule, we pick the nonbasic variable with smallest index,  $x_1$ . There are two limitations to the growth of  $x_1$ , i.e., given by  $y_1 = 1 - 2x_1$  and  $y_2 = 5 - 3x_1$ . Therefore, the tightest limit is  $\min\{\frac{1}{2}, \frac{5}{3}\} = \frac{1}{2}$ , given by the basic variable  $y_1$ , which leaves the basis. Pivoting is performed on coefficient 2 in tableau position(1,1)

1. divide row 1 by 2
2. add 5 times row 1 to row 0
3. subtract 3 times row 1 from row 2

We obtain the tableau

		$x_1$	$x_2$	$x_3$	$y_1$	$y_2$
	$-\frac{7}{2}$	0	-2	$\frac{11}{2}$	$\frac{5}{2}$	0
$x_1$	$\frac{1}{2}$	1	0	$\frac{3}{2}$	$\frac{1}{2}$	0
$y_2$	$\frac{7}{2}$	0	2	$-\frac{11}{2}$	$-\frac{3}{2}$	1

which shows the current basic solution  $\underline{x}_B = (x_1, y_2)^T = (1/2, 7/2)^T$  of value  $7/2$  (obviously  $\underline{x}_N = 0$ ). Since the only negative reduced cost is given by  $x_2$ ,  $x_2$  enters the basis. The only limit to its growth is given by  $y_2 = \frac{7}{2} - 2x_2$ , therefore  $x_2 \leq \frac{7}{4}$  and  $y_2$  leaves the basis. Pivoting is performed on coefficient 2 in position (2,2) of the previous tableau

1. add row 2 to row 0
2. divide row 2 by 2

We obtain

	$x_1$	$x_2$	$x_3$	$y_1$	$y_2$
	0	0	0	1	1
$x_1$	$\frac{1}{2}$	1	0	$\frac{3}{2}$	$\frac{1}{2}$
$x_2$	$\frac{7}{4}$	0	1	$-\frac{11}{4}$	$-\frac{3}{4}$

Since the reduced costs for the nonbasic variables are all nonnegative, Phase I stops, proving that the feasible region of the original problem is nonempty, and yielding an initial basic solution of value  $\underline{x}_B = (x_1, x_2)$ .

#### PHASE II

The objective function is  $x_1 - 2x_2$ , and the initial basis is  $(x_1, x_2)$ , with  $x_3$  nonbasic. We express  $x_1, x_2$  w.r.t.  $x_3$

$$\begin{aligned} x_1 &= \frac{1}{2} - \frac{3}{2}x_3 \\ x_2 &= \frac{7}{4} + \frac{11}{4}x_3 \end{aligned}$$

The objective function becomes  $-3 - 7x_3$ . Removing from the tableau the columns of the auxiliary variables, we obtain

	$x_1$	$x_2$	$x_3$
	3	0	-7
$x_1$	$\frac{1}{2}$	1	$\frac{3}{2}$
$x_2$	$\frac{7}{4}$	0	$-\frac{11}{4}$

Variable  $x_3$ , having a negative reduced cost of -7, enters the basis. Since the only limit to its growth is given by  $x_1 = \frac{1}{2} - \frac{3}{2}x_3$ ,  $x_1$  leaves the basis. Therefore, we perform a pivoting operation on coefficient  $\frac{3}{2}$  in position (3,1)

1. divide row 1 by  $\frac{3}{2}$ ;
2. add 7 times row 1 to row 0
3. add  $-\frac{11}{4}$  times row 1 to row 3

We obtain

	$x_1$	$x_2$	$x_3$
	$\frac{16}{3}$	$\frac{14}{3}$	0
$x_3$	$\frac{1}{3}$	$\frac{2}{3}$	1
$x_2$	$\frac{8}{3}$	$\frac{11}{6}$	0

The reduced costs are all nonnegative and the algorithm halts, yielding the optimal solution  $(0, \frac{8}{3}, \frac{1}{3})$ , of value  $z^* = -\frac{16}{3}$ .