

Exercises on ILP formulations

1. Given the following set S of integer solutions:

$S = \{(0,0,0,0), (1,0,0,0), (0,1,0,0), (0,0,1,0), (0,0,0,1), (0,1,0,1), (0,0,1,1)\}$ and the two polyhedron:

$$P_1 = \{x \in \mathbb{R}^4 : 0 \leq x \leq 1, 83x_1 + 61x_2 + 49x_3 + 20x_4 \leq 100\}$$

$$P_2 = \{x \in \mathbb{R}^4 : 0 \leq x \leq 1, 4x_1 + 3x_2 + 2x_3 + x_4 \leq 4\}$$

- verify that both P_1 and P_2 are formulations for S ;
- establish which of the two formulations is the best one.

2. Consider a transport problem with m possible sources (plants) and n destinations (customers). In many applications, the problem of determining which of the possible origins must work arises, since opening a source i generates a startup fixed cost F_i . Are also known costs c_{ij} to transport a single product from the source i to the destination j and the demand d_j of customer j . The aim is to determine the opening strategy of the plants and the transport plan with minimum total cost.

Let us introduce the variables $x_{ij} \geq 0$ to represent the quantity transported from origin i to destination j and the binary variables y_i such that:

$$y_i = \begin{cases} 1 & \text{if plant } i \text{ is active} \\ 0 & \text{otherwise} \end{cases}$$

The problem can be modeled as P_1 :

$$\min \sum_{i=1}^m \sum_{j=1}^n c_{ij} x_{ij} + \sum_{i=1}^m F_i y_i$$

$$\sum_{i=1}^m x_{ij} = d_j \quad \text{for } j=1, \dots, n \quad (3.1)$$

$$\sum_{j=1}^n x_{ij} \leq D y_i \quad \text{for } i=1, \dots, m \quad (3.2)$$

$$x_{ij} \geq 0 \quad \text{for } i=1, \dots, m, \text{ for } j=1, \dots, n$$

$$y_i \in \{0,1\} \quad \text{for } i=1, \dots, m$$

with $D = \sum_{j=1}^n d_j$.

Another possible formulation is P_2 that differs from P_1 only in constraints (3.2) that are replaced with the following mn constraints:

$$x_{ij} \leq d_j y_i \quad \text{for } j = 1, \dots, n \quad (3.3)$$

State and prove which of the two formulations is better.