

# Sharpening through spatial filtering

Stefano Ferrari

Università degli Studi di Milano  
stefano.ferrari@unimi.it

## Methods for Image Processing

academic year 2018–2019

### *Sharpening*

- ▶ The term *sharpening* is referred to the techniques suited for enhancing the intensity transitions.
- ▶ In images, the borders between objects are perceived because of the intensity change: the crisper the intensity transitions, the sharper the image is perceived.
- ▶ The intensity transition between adjacent pixels is related to the derivatives of the image in that position.
- ▶ Hence, operators (possibly expressed as linear filters) able to compute the derivatives of a digital image are very interesting.

## First derivative of an image

- ▶ Since the image is a discrete function, the traditional definition of derivative cannot be applied.
- ▶ Hence, a suitable operator have to be defined such that it satisfies the main properties of the first derivative:
  1. equal to zero in the regions where the intensity is constant;
  2. different from zero for an intensity transition;
  3. constant on ramps where the intensity transition is constant.
- ▶ The natural derivative operator is the difference between the intensity of neighboring pixels (spatial differentiation).
- ▶ For simplicity, the monodimensional case can be considered:

$$\frac{\partial f}{\partial x} = f(x + 1) - f(x)$$

- ▶ Since  $\frac{\partial f}{\partial x}$  is defined using the next pixel:
  - ▶ it cannot be computed for the last pixel of each row (and column);
  - ▶ it is different from zero in the pixel before a step.


## Second derivative of an image

- ▶ Similarly, the second derivative operator can be defined as:

$$\begin{aligned}\frac{\partial^2 f}{\partial x^2} &= f(x + 1) - f(x) - (f(x) - f(x - 1)) \\ &= f(x + 1) - 2f(x) + f(x - 1)\end{aligned}$$

- ▶ This operator satisfies the following properties:
  1. it is equal to zero where the intensity is constant;
  2. it is different from zero at the beginning of a step (or a ramp) of the intensity;
  3. it is equal to zero on the constant slope ramps.
- ▶ Since  $\frac{\partial^2 f}{\partial x^2}$  is defined using the previous and the next pixels:
  - ▶ it cannot be computed with respect to the first and the last pixels of each row (and column);
  - ▶ it is different from zero in the pixel that precedes and in the one that follows a step.

## Derivatives of an image: an example



Note: at the step, the second derivative's sign switches (*zero crossing*).

## Laplacian

- ▶ Usually the sharpening filters make use of the second order operators.
  - ▶ A second order operator is more sensitive to intensity variations than a first order operator.
- ▶ Besides, partial derivatives has to be considered for images.
  - ▶ The derivative in a point depends on the direction along which it is computed.
- ▶ Operators that are invariant to rotation are called *isotropic*.
  - ▶ Rotate and differentiate (or filtering) has the same effects of differentiate and rotate.
- ▶ The *Laplacian* is the simplest isotropic derivative operator (wrt. the principal directions):

$$\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2}$$

## Laplacian filter

- In a digital image, the second derivatives wrt.  $x$  and  $y$  are computed as:

$$\frac{\partial^2 f}{\partial x^2} = f(x+1, y) - 2f(x, y) + f(x-1, y)$$
$$\frac{\partial^2 f}{\partial y^2} = f(x, y+1) - 2f(x, y) + f(x, y-1)$$

- Hence, the Laplacian results:

$$\nabla^2 f(x, y) = f(x+1, y) + f(x-1, y) + f(x, y+1) + f(x, y-1) - 4f(x, y)$$

- Also the derivatives along to the diagonals can be considered:

$$\begin{aligned} \nabla^2 f(x, y) &+ f(x-1, y-1) + f(x+1, y+1) \\ &+ f(x-1, y+1) + f(x+1, y-1) - 4f(x, y) \end{aligned}$$

## Laplacian filter (2)

0	1	0
1	-4	1
0	1	0


Laplacian filter invariant to  $90^\circ$  rotations

1	1	1
1	-8	1
1	1	1

Laplacian filter invariant to  $45^\circ$  rotations

## Laplacian filter: example

- ▶ The Laplacian has often negative values.
- ▶ In order to be visualized, it must be properly scaled to the representation interval  $[0, \dots, L - 1]$ .



(a)

(b)

(c)

(a) Original image, (b) its Laplacian, (c) its Laplacian scaled such that zero is displayed as the intermediate gray level.

## Laplacian filter: example (2)

- ▶ The Laplacian is positive at the onset of a step and negative at the end.
- ▶ Subtracting the Laplacian (or a fraction of it) from the image, the height of the step is increased:  $g = f + c\nabla^2 f$ ,  $-1 \leq c \leq 0$



(a)

(b)

(c)

(a) Original image, (b) Laplacian filtered, (c) Laplacian with diagonals filtered.

## Unsharp masking


- ▶ The technique known as *unsharp masking* is a method of common use in graphics for making the images sharper.
- ▶ It consists of:
  1. defocusing the original image;
  2. obtaining the mask as the difference between the original image and its defocused copy;
  3. adding the mask to the original image.
- ▶ The process can be formalized as:

$$g = f + k \cdot (f - f * h)$$

where  $f$  is the original image,  $h$  is the smoothing filter and  $k$  is a constant for tuning the mask contribution.

- ▶ If  $k > 1$ , the process is called *highboost* filtering.

## Unsharp masking (2)



## Gradient

- ▶ The *gradient* of a function is the vector formed by its partial derivatives.
- ▶ For a bidimensional function,  $f(x, y)$ :

$$\nabla f \equiv \text{grad}(f) \equiv \begin{bmatrix} g_x \\ g_y \end{bmatrix} = \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \end{bmatrix}$$

- ▶ The gradient vector points toward the direction of maximum variation.
- ▶ The gradient *magnitude*,  $M(x, y)$  is:

$$M(x, y) = \text{mag}(\nabla f) = \sqrt{g_x^2 + g_y^2}$$

- ▶ It is also called *gradient image*.
- ▶ Often approximated as  $M(x, y) \approx |g_x| + |g_y|$ .

## Derivative operators

- ▶ Basic definitions:

$$g_x(x, y) = f(x + 1, y) - f(x, y)$$

$$g_y(x, y) = f(x, y + 1) - f(x, y)$$

$$g_x: \begin{bmatrix} -1 & 1 \end{bmatrix} \quad g_y: \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- ▶ Roberts operators:

$$g_x(x, y) = f(x + 1, y + 1) - f(x, y)$$

$$g_y(x, y) = f(x, y + 1) - f(x - 1, y)$$

$$g_x: \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \quad g_y: \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

## Derivative operators (2)

- Sobel operators:

$$\begin{aligned}g_x(x, y) = & -f(x-1, y-1) - 2f(x-1, y) \\ & - f(x-1, y+1) + f(x+1, y-1) \\ & + 2f(x+1, y) + f(x+1, y+1)\end{aligned}$$

$$\begin{aligned}g_y(x, y) = & -f(x-1, y-1) - 2f(x, y-1) \\ & - f(x+1, y-1) + f(x-1, y+1) \\ & + 2f(x, y+1) + f(x+1, y+1)\end{aligned}$$


$g_x$ :

-1	-2	-1
0	0	0
1	2	1

$g_y$ :

-1	0	1
-2	0	2
-1	0	1

## Example of gradient based application




- The Sobel filtering reduces the visibility of those regions in which the intensity changes slowly, allowing to highlight the defects (and making defects detection easier for automatic processing).




## Combined methods

- ▶ Often, a single technique is not sufficient for obtaining the desired results.
- ▶ For example, the image (a) is affected by noise and has a narrow dynamic range.



(a)




(b)


- ▶ Laplacian filtering (b) enhance the details, but also the noise.
- ▶ The gradient is less sensitive to the noise than the Laplacian (which is a second order operator).

## Combined methods (2)

- ▶ The gradient, smoothed in order to avoid the noise, can be used for weighting the contribution of the Laplacian.
- ▶ The gradient smoothed using a  $5 \times 5$  averaging filter is reported in (c); Sobel filters have been used for the gradient.



(c)




(d)


- ▶ This mask multiplied by the Laplacian results in the image in (d).
- ▶ The intensity changes are preserved, while the noise has been attenuated.

### Combined methods (3)

- ▶ The image (d) can be added to the original image, which results in the image (e).
- ▶ The dynamic range can be enlarged applying a power transformation (f).



(e)



(f)

- ▶ The intensity transformation make the noise more visible, but also enhance other details, such as the tissues around the skeleton.

### Bilateral Filtering \*

The image  $g$  is obtained from  $f$ , through bilateral filtering:

$$g(p) = \frac{1}{W_p} \sum_{q \in N_p} \exp\left(-\frac{\|q - p\|^2}{\sigma_s^2}\right) \exp\left(-\frac{\|f(q) - f(p)\|^2}{\sigma_i^2}\right) f(q)$$


where  $W_p$  is the normalization factor:

$$W_p = \sum_{q \in N_p} \exp\left(-\frac{\|q - p\|^2}{\sigma_s^2}\right) \exp\left(-\frac{\|f(q) - f(p)\|^2}{\sigma_i^2}\right)$$

and  $N_p$  is a suitable neighborhood of  $p$ .


- ▶ What is the effect produced by the filter?
- ▶ Notes:
  - ▶ when  $\sigma_i$  grows, the filter tends to an averaging filter;
  - ▶ the filter is not linear.

## Bilateral Filtering \*




- (a) original image ( $100 \times 100$ , 256 gray levels)
- (b) after filtering with Gaussian filter ( $7 \times 7$ ,  $\sigma_s = 3$ )
- (c) after filtering with bilateral filter ( $7 \times 7$ ,  $\sigma_s = 3$ ,  $\sigma_i = 50$ )

## Bilateral Filtering \*



- (a) Original image ( $107 \times 90$ , 256 gray levels)
- (b) after filtering with Gaussian filter ( $7 \times 7$ ,  $\sigma_s = 3$ )
- (c) after filtering with bilateral filter ( $7 \times 7$ ,  $\sigma_s = 3$ ,  $\sigma_i = 30$ )

## Homeworks and suggested readings



DIP, Sections 3.6, 3.7

- ▶ pp. 157–173



GIMP

- ▶ Filters
  - ▶ Enhance
    - ▶ Sharpen
    - ▶ Unsharp mask
  - ▶ Generic
    - ▶ Convolution matrix