
Università degli Studi di Milano
Master Degree in Computer Science

Information Management course

Teacher: Alberto Ceselli

Lecture 02 : 03/10/2012


Data Mining: Concepts and Techniques

— Chapter 2 —

Jiawei Han, Micheline Kamber, and Jian Pei
University of Illinois at Urbana-Champaign
Simon Fraser University

©2012 Han, Kamber, and Pei. All rights reserved.

Chapter 2: Getting to Know Your Data

- Data Objects and Attribute Types 
- Basic Statistical Descriptions of Data
- Data Visualization
- Measuring Data Similarity and Dissimilarity
- Summary

Types of Data Sets

- Record
 - Relational records
 - Data matrix, e.g., numerical matrix, crosstabs
 - Document data: text documents: term-frequency vector
 - Transaction data
- Graph and network
 - World Wide Web
 - Social or information networks
 - Molecular Structures
- Ordered
 - Video data: sequence of images
 - Temporal data: time-series
 - Sequential Data: transaction sequences
 - Genetic sequence data
- Spatial, image and multimedia:
 - Spatial data: maps
 - Image data: .bmp
 - Video data: .avi

<i>TID</i>	<i>Items</i>
1	Bread, Coke, Milk
2	Beer, Bread
3	Beer, Coke, Diaper, Milk
4	Beer, Bread, Diaper, Milk
5	Coke, Diaper, Milk

Important Characteristics of Structured Data

- Dimensionality
 - Curse of dimensionality
(the volume of the space grows fast with the number of dimensions, and the available data becomes sparse)
- Sparsity
 - Only presence counts
- Resolution
 - Patterns depend on the scale
- Distribution
 - Centrality and dispersion

Data Objects

- Data sets are made up of data objects.
- A **data object** represents an entity.
- Examples:
 - sales database: customers, store items, sales
 - medical database: patients, treatments
 - university database: students, professors, courses
- Also called *samples*, *examples*, *instances*, *data points*, *objects*, *tuples*.
- Data objects are described by **attributes**.
- Database rows -> data objects; columns -> attributes.

Attributes

- **Attribute (or dimensions, features, variables)**: a data field, representing a characteristic or feature of a data object.
 - *E.g., customer_ID, name, address*
- Types:
 - Nominal
 - Binary
 - Ordinal
 - Numeric: quantitative
 - Interval-scaled
 - Ratio-scaled

Attribute Types

- **Nominal:** categories, states, or “names of things”
 - *Hair_color* = {*auburn, black, blond, brown, grey, red, white*}
 - marital status, occupation, ID numbers, zip codes
- **Binary**
 - Nominal attribute with only 2 states (0 and 1)
 - Symmetric binary: both outcomes equally important
 - e.g., gender
 - Asymmetric binary: outcomes not equally important.
 - e.g., medical test (positive vs. negative)
 - Convention: assign 1 to most important outcome (e.g., HIV positive)
- **Ordinal**
 - Values have a meaningful order (ranking) but magnitude between successive values is not known.
 - *Size* = {*small, medium, large*}, grades, army rankings

Numeric Attribute Types

- Quantity (integer or real-valued)
- **Interval**
 - Measured on a scale of **equal-sized units**
 - Values have order
 - E.g., *temperature in C° or F°, calendar dates*
 - No true zero-point
- **Ratio**
 - Inherent **zero-point**
 - We can speak of values as being an order of magnitude larger than the unit of measurement (10 K° is twice as high as 5 K°).
 - e.g., *temperature in Kelvin, length, counts, monetary quantities*

Discrete vs. Continuous Attributes (ML view)


■ Discrete Attribute

- Has only a finite or countably infinite set of values
 - E.g., zip codes, profession, or the set of words in a collection of documents
- Sometimes, represented as integer variables
- Note: Binary attributes are a special case of discrete attributes

■ Continuous Attribute

- Has real numbers as attribute values
 - E.g., temperature, height, or weight
- Practically, real values can only be measured and represented using a finite number of digits
- Continuous attributes are typically represented as floating-point variables

Chapter 2: Getting to Know Your Data

- Data Objects and Attribute Types
- Basic Statistical Descriptions of Data 
- Data Visualization
- Measuring Data Similarity and Dissimilarity
- Summary

Basic Statistical Descriptions of Data

- Motivation
 - To better understand the data: central tendency, variation and spread
- Data dispersion characteristics
 - median, max, min, quantiles, outliers, variance...
- Numerical dimensions correspond to sorted intervals
 - Data dispersion: analyzed with multiple granularities of precision
 - Boxplot or quantile analysis on sorted intervals
- Dispersion analysis on computed measures
 - Folding measures into numerical dimensions
 - Boxplot or quantile analysis on the transformed cube

Measuring the Central Tendency

- Mean (algebraic measure) (sample vs. population):

Note: n is sample size and N is population size.

- Weighted arithmetic mean
- Sensitive to outliers: trimmed mean (chopping extreme values)

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

$$\bar{x} = \frac{\sum_{i=1}^n w_i x_i}{\sum_{i=1}^n w_i}$$

- Median:

- Middle value if odd number of values, or average of the middle two values otherwise
- Estimated by interpolation (for *grouped data*):

age	frequency
1–5	200
6–15	450
16–20	300
21–50	1500
51–80	700
81–110	44

$$\text{median} = L_1 + \left(\frac{\frac{n}{2} - (\sum \text{freq})_l}{\text{freq}_{\text{median}}} \right) \text{width}$$

Lower boundary of the median interval

values in the dataset

Freq. of the median interval

Sum of freq. of intervals preceding the median

Measuring the Central Tendency

■ Mode

- Value that occurs most frequently in the data
- Unimodal, bimodal, trimodal
- Empirical formula for moderately skewed:

$$mean - mode \simeq 3 \times (mean - median)$$

Mean: 58

Median: $(52+56)/2 = 54$




Mode: 52 and 70 (bimodal)

Midrange: $(30+110) / 2 = 70$

Employed	Salary
1	30
2	36
3	47
4	50
5	52
6	52
7	56
8	60
9	63
10	70
11	70
12	110

Symmetric vs. Skewed Data

- Median, mean and mode of symmetric, positively and negatively skewed data



Measuring the Dispersion of Data

- Quartiles, outliers and boxplots
 - **Quartiles:** Q_1 (25th percentile), Q_3 (75th percentile)
 - **Inter-quartile range:** $IQR = Q_3 - Q_1$
 - **Five number summary:** min, Q_1 , median, Q_3 , max (nice for skewed distributions)
 - **Boxplot:** ends of the box are the quartiles; median is marked; add whiskers, and plot outliers individually
 - **Outlier:** usually, a value higher/lower than $1.5 \times IQR$
- Variance and standard deviation (*sample: s , population: σ*)
 - **Variance:** (algebraic, scalable computation)

$$s^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2 = \frac{1}{n} \left[\sum_{i=1}^n x_i^2 - \frac{1}{n} \left(\sum_{i=1}^n x_i \right)^2 \right]$$

$$\sigma^2 = \frac{1}{N} \sum_{i=1}^n (x_i - \mu)^2 = \frac{1}{N} \sum_{i=1}^n x_i^2 - \mu^2$$

- **Standard deviation s (or σ)** is the square root of variance



Boxplot Analysis

- **Five-number summary** of a distribution


- Minimum, Q1, Median, Q3, Maximum

- **Boxplot**

- Data is represented with a box
- The ends of the box are at the first and third quartiles, i.e., the height of the box is IQR
- The median is marked by a line within the box
- Whiskers: two lines outside the box extended to Minimum and Maximum
- Outliers: points beyond a specified outlier threshold, plotted individually




Visualization of Data Dispersion: 3-D Boxplots



Properties of Normal Distribution Curve

- The normal (distribution) curve
 - From $\mu - \sigma$ to $\mu + \sigma$: contains about 68% of the measurements (μ : mean, σ : standard deviation)
 - From $\mu - 2\sigma$ to $\mu + 2\sigma$: contains about 95% of it
 - From $\mu - 3\sigma$ to $\mu + 3\sigma$: contains about 99.7% of it



Graphic Displays of Basic Statistical Descriptions



- **Boxplot:** graphic display of five-number summary
- **Histogram:** x-axis are values, y-axis repres. frequencies
- **Quantile plot:** each value x_i is paired with f_i indicating that approximately $100 f_i\%$ of data are $\leq x_i$
- **Quantile-quantile (q-q) plot:** graphs the quantiles of one univariant distribution against the corresponding quantiles of another
- **Scatter plot:** each pair of values is a pair of coordinates and plotted as points in the plane

Histogram Analysis

- Histogram: Graph display of tabulated frequencies, shown as bars
- It shows what proportion of cases fall into each of several categories
- Differs from a bar chart in that it is the *area* of the bar that denotes the value, not the height as in bar charts, a crucial distinction when the categories are not of uniform width
- The categories are usually specified as non-overlapping intervals of some variable. The categories (bars) must be adjacent

40


Histograms Often Tell More than Boxplots



- The two histograms shown in the left may have the same boxplot representation
 - The same values for: min, Q1, median, Q3, max
- But they have rather different data distributions


Quantile Plot

- Displays all of the data (allowing the user to assess both the overall behavior and unusual occurrences)
- Plots **quantile** information
 - For a data x_i data sorted in increasing order, f_i indicates that approximately 100 f_i % of the data are below or equal to the value x_i




Quantile-Quantile (Q-Q) Plot

- Graphs the quantiles of one univariate distribution against the corresponding quantiles of another
- View: Is there is a shift in going from one distribution to another?
- Example shows unit price of items sold at Branch 1 vs. Branch 2 for each quantile. Unit prices of items sold at Branch 1 tend to be lower than those at Branch 2.






Scatter plot

- Provides a first look at bivariate data to see clusters of points, outliers, etc
- Each pair of values is treated as a pair of coordinates and plotted as points in the plane






Positively and Negatively Correlated Data




- The left half fragment is positively correlated
- The right half is negative correlated

Uncorrelated Data



Chapter 2: Getting to Know Your Data

- Data Objects and Attribute Types
- Basic Statistical Descriptions of Data
- Data Visualization
- Measuring Data Similarity and Dissimilarity 
- Summary

Similarity and Dissimilarity

- **Similarity**
 - Numerical measure of how alike two data objects are
 - Value is higher when objects are more alike
 - Often falls in the range $[0,1]$
- **Dissimilarity** (e.g., distance)
 - Numerical measure of how different two data objects are
 - Lower when objects are more alike
 - Minimum dissimilarity is often 0
 - Upper limit varies
- **Proximity** refers to a similarity or dissimilarity

Proximity Measures for Binary Attributes

- A contingency table for binary data

		Object j		
		1	0	sum
Object i	1	q	r	$q+r$
	0	s	t	$s+t$
sum		$q+s$	$r+t$	p

- Distance measure for symmetric bin. vars (0 and 1 equally important):

$$d(i, j) = \frac{r + s}{q + r + s + t}$$

- Distance measure for asymm. bin. vars (1 more important - e.g. diseases):

$$d(i, j) = \frac{r + s}{q + r + s}$$

- Jaccard coefficient (*similarity* measure for *asymmetric* binary variables):

$$sim_{Jaccard}(i, j) = \frac{q}{q + r + s}$$

- Note: Jaccard coefficient is the same as “coherence”:

$$coherence(i, j) = \frac{sup(i, j)}{sup(i) + sup(j) - sup(i, j)} = \frac{q}{(q + r) + (q + s) - q}$$

Dissimilarity between Binary Variables

- Example

Name	Gender	Fever	Cough	Test-1	Test-2	Test-3	Test-4
Jack	M	Y	N	P	N	N	N
Mary	F	Y	N	P	N	P	N
Jim	M	Y	P	N	N	N	N

- Gender is a symmetric attribute (let's discard it!)
- The remaining attributes are asymmetric binary
- Let the values Y and P be 1, and the value N 0

$$d(\text{jack}, \text{mary}) = \frac{0 + 1}{2 + 0 + 1} = 0.33$$

$$d(\text{jack}, \text{jim}) = \frac{1 + 1}{1 + 1 + 1} = 0.67$$

$$d(\text{jim}, \text{mary}) = \frac{1 + 2}{1 + 1 + 2} = 0.75$$

Proximity Measures for Nominal Attributes

- Can take 2 or more states, e.g., red, yellow, blue, green (generalization of a binary attribute)

- Method 1: Simple matching

- m : # of matches, p : total # of variables

$$d(i, j) = \frac{p - m}{p}$$

- Method 2: Use a large number of binary attributes

- creating a new binary attribute for each of the M nominal states

Proximity on Numeric Data: Minkowski Distance

- *Minkowski distance*: A popular distance measure

$$d(i, j) = \sqrt[h]{|x_{i1} - x_{j1}|^h + |x_{i2} - x_{j2}|^h + \dots + |x_{ip} - x_{jp}|^h}$$

where $i = (x_{i1}, x_{i2}, \dots, x_{ip})$ and $j = (x_{j1}, x_{j2}, \dots, x_{jp})$ are two p -dimensional data objects, and h is the order (the distance so defined is also called L- h norm)

- Properties
 - $d(i, j) > 0$ if $i \neq j$, and $d(i, i) = 0$ (Positive definiteness)
 - $d(i, j) = d(j, i)$ (Symmetry)
 - $d(i, j) \leq d(i, k) + d(k, j)$ (Triangle Inequality)
- A distance that satisfies these properties is a **metric**

Special Cases of Minkowski Distance

- $h = 1$: **Manhattan** (city block, L_1 norm) **distance**
 - E.g., the Hamming distance: the number of bits that are different between two binary vectors

$$d(i, j) = |x_{i1} - x_{j1}| + |x_{i2} - x_{j2}| + \dots + |x_{ip} - x_{jp}|$$

- $h = 2$: (L_2 norm) **Euclidean** distance

$$d(i, j) = \sqrt{(|x_{i1} - x_{j1}|^2 + |x_{i2} - x_{j2}|^2 + \dots + |x_{ip} - x_{jp}|^2)}$$

- $h \rightarrow \infty$. **“supremum”** (L_{\max} norm, L_{∞} norm) distance.
 - This is the maximum difference between any component (attribute) of the vectors

$$d(i, j) = \lim_{h \rightarrow \infty} \left(\sum_{f=1}^p |x_{if} - x_{jf}|^h \right)^{\frac{1}{h}} = \max_f |x_{if} - x_{jf}|$$

Example: Minkowski Distance

point I

Example (L_2)

Supremum (L_∞)

Standardizing Numeric Data

- Z-score: $z = \frac{x - \mu}{\sigma}$
 - X: raw data, μ : mean of the population, σ : standard deviation
 - the distance between the raw score and the population mean in units of the standard deviation
 - <0 when the raw score is below the mean, >0 when above
- An alternative way: Calculate the mean absolute deviation

$$s_f = \frac{1}{n} (|x_{1f} - m_f| + |x_{2f} - m_f| + \dots + |x_{nf} - m_f|)$$

where

$$m_f = \frac{1}{n} (x_{1f} + x_{2f} + \dots + x_{nf}).$$

- standardized measure (z-score): $z_{if} = \frac{x_{if} - m_f}{s_f}$
- mean absolute deviation is more robust than std dev

Ordinal Variables

- An ordinal variable can be discrete or continuous
- Order is important, e.g., rank
- Can be treated like interval-scaled
 - replace x_{if} by their rank $r_{if} \in \{1, \dots, M_f\}$
 - map (normalize) the range of each variable onto $[0, 1]$ by replacing x_{if} by

$$z_{if} = \frac{r_{if} - 1}{M_f - 1}$$

- compute the dissimilarity using distance measures for numeric attributes

Attributes of Mixed Type

- A database may contain all attribute types
 - Nominal, symmetric binary, asymmetric binary, numeric, ordinal
- One may use a weighted formula to combine their effects

$$d(i, j) = \frac{\sum_{f=1}^p \delta_{ij}^{(f)} d_{ij}^{(f)}}{\sum_{f=1}^p \delta_{ij}^{(f)}}$$

- Choice of $\delta_{ij}^{(f)}$
 - Set $\delta_{ij}^{(f)} = 0$ if
 - x_{if} or x_{jf} is missing
 - $x_{if} = x_{jf} = 0$ and f is asymmetric binary
 - Set $\delta_{ij}^{(f)} = 1$ otherwise

Attributes of Mixed Type

$$d(i, j) = \frac{\sum_{f=1}^p \delta_{ij}^{(f)} d_{ij}^{(f)}}{\sum_{f=1}^p \delta_{ij}^{(f)}}$$

- Choice of $d_{ij}^{(f)}$
 - when f is binary or nominal:
 $d_{ij}^{(f)} = 0$ if $x_{if} = x_{jf}$, $d_{ij}^{(f)} = 1$ otherwise
 - when f is numeric: use the normalized distance
 - when f is ordinal
 - Compute ranks r_{if} and $z_{if} = \frac{r_{if} - 1}{M_f - 1}$
 - Treat z_{if} as interval-scaled

Cosine Similarity

- A **document** can be represented by thousands of attributes, each recording the *frequency* of a particular word (such as keywords) or phrase in the document.

<i>Document</i>	<i>teamcoach</i>	<i>hockey</i>	<i>baseball</i>	<i>soccer</i>	<i>penalty</i>	<i>score</i>	<i>win</i>	<i>loss</i>	<i>season</i>
Document1	5	0	3	0	2	0	2	0	0
Document2	3	0	2	0	1	1	1	0	1
Document3	0	7	0	2	1	0	3	0	0
Document4	0	1	0	0	1	2	0	3	0

- Other vector objects: gene features in micro-arrays, ...
- Applications: information retrieval, biologic taxonomy, gene feature mapping, ...
- Issue: very long and **sparse**
- Treat documents as vectors, and compute a **cosine similarity**

Cosine Similarity

- Cosine measure: If x and y are two vectors (e.g., term-frequency vectors), then

$$\cos(x, y) = (x \bullet y) / \|x\| \|y\|$$

where

- \bullet indicates vector dot product,
- $\|x\|$: the L2 norm (length) of vector x $\|x\| = \sqrt{x_1^2 + x_2^2 + \dots + x_p^2}$
- Remark: when attributes are binary valued:
 - \bullet indicates the number of shared features
 - $\|x\| \|y\|$ is the geometric mean between the number of features of x and the number of features of y :
$$\text{sqrt}(a) * \text{sqrt}(b) = \text{sqrt}(a * b)$$
 - $\cos(x, y)$ measures relative possession of common features

Example: Cosine Similarity

- $\cos(x, y) = (x \bullet y) / (\|x\| \|y\|)$
- Ex: Find the **similarity** between documents x and y.

$$\mathbf{x} = (5, 0, 3, 0, 2, 0, 0, 2, 0, 0)$$

$$\mathbf{y} = (3, 0, 2, 0, 1, 1, 0, 1, 0, 1)$$

$$\begin{aligned} x \bullet y &= 5*3+0*0+3*2+0*0+2*1+0*1+0*1+2*1+0*0+0*1= \\ &= 25 \end{aligned}$$

$$\begin{aligned} \|x\| &= (5*5+0*0+3*3+0*0+2*2+0*0+0*0+2*2+0*0+0*0)^{0.5}= \\ &= 6.481 \end{aligned}$$

$$\begin{aligned} \|y\| &= (3*3+0*0+2*2+0*0+1*1+1*1+0*0+1*1+0*0+1*1)^{0.5}= \\ &= 4.12 \end{aligned}$$

$$\cos(x, y) = 25 / (6.481 * 4.12) = 0.94$$

References

- W. Cleveland, Visualizing Data, Hobart Press, 1993
- T. Dasu and T. Johnson. Exploratory Data Mining and Data Cleaning. John Wiley, 2003
- U. Fayyad, G. Grinstein, and A. Wierse. Information Visualization in Data Mining and Knowledge Discovery, Morgan Kaufmann, 2001
- L. Kaufman and P. J. Rousseeuw. Finding Groups in Data: an Introduction to Cluster Analysis. John Wiley & Sons, 1990.
- H. V. Jagadish et al., Special Issue on Data Reduction Techniques. Bulletin of the Tech. Committee on Data Eng., 20(4), Dec. 1997
- D. A. Keim. Information visualization and visual data mining, IEEE trans. on Visualization and Computer Graphics, 8(1), 2002
- D. Pyle. Data Preparation for Data Mining. Morgan Kaufmann, 1999
- S. Santini and R. Jain, " Similarity measures", IEEE Trans. on Pattern Analysis and Machine Intelligence, 21(9), 1999
- E. R. Tufte. The Visual Display of Quantitative Information, 2nd ed., Graphics Press, 2001
- C. Yu et al., Visual data mining of multimedia data for social and behavioral studies, Information Visualization, 8(1), 2009